arXiv:1510.00692v1 [math.OA] 2 Oct 2015

FAITHFUL ACTIONS OF LOCALLY COMPACT QUANTUM GROUPS ON CLASSICAL SPACES

DEBASHISH GOSWAMI AND SUTANU ROY

Abstract. It is well-known that no non-Kac compact quantum group can faithfully act on $\mathrm{C}(X)$ for a classical, compact Hausdorff space X. However, in this article we show that this is no longer true if we go to non-compact spaces and non-compact quantum groups, by exhibiting a large class of examples of locally compact quantum groups coming from bicrossed product construction, including non-Kac ones, which can faithfully and ergodically act on classical (non-compact) spaces. However, none of these actions can be isometric in the sense of [5], leading to the conjecture that the result obtained by Goswami and Joardar in [6] about non-existence of genuine quantum isometry of classical compact connected Riemannian manifolds may hold in the non-compact case as well.

1. Introduction

Symmetry plays a crucial role in many areas of mathematics and physics. Conventionally, group actions are used to model symmetries and with the advent of more general mathematical structures called quantum groups in [3, 9, 17]. It should be natural to consider the actions (defined in a suitable sense depending on the algebraic or analytic framework chosen) of quantum groups on classical and non-commutative spaces. In this context, a very interesting programme is to study quantum symmetries of classical spaces. One may hope that there are many more quantum symmetries of a given classical space than classical group symmetries which will help one understand the space better. Indeed, it has been a remarkable discovery of S. Wang that for $n \geq 4$, a finite set of cardinality n has an infinite dimensional compact quantum group ('quantum permutation group') of symmetries. For the relevance of quantum group symmetries in a wider and more geometric context, we refer the reader to the discussion on 'hidden symmetry in algebraic geometry' in [12, Chapter 13] where Manin made a remark about possible genuine Hopf algebra symmetries of classical smooth algebraic varieties.

Recently, several examples of faithful continuous action by genuine (i.e. not of the form $\mathrm{C}(G)$ for a compact group G) compact quantum groups on $\mathrm{C}(X)$ for a connected compact space X were constructed by H. Huang [7]. In [4] an example of a faithful action by the finite dimensional genuine compact quantum group on the algebra of regular function of a non-smooth variety was given. However, it turns out rather formidable to construct such actions when the space is smooth (and connected) and the action is also smooth in some natural sense. In [6], it is conjectured that no smooth faithful action of genuine compact quantum group on a compact connected smooth manifold can exist. The conjecture has been proved in

2010 *Mathematics Subject Classification.* 81R50, 46L89.

Key words and phrases. Bicrossed product, faithful C*-action, locally compact quantum group.

The first author was partially supported by Swarnajayanti grant given by D.S.T., Government of India. The second author was supported by the visiting scientist program at the Indian Statistical Institute, Kolkata. Parts of the manuscript had been written and revised while the second author had been supported by Fields–Ontario postdoctoral fellowship, NSERC and ERA at the University of Ottawa and Carleton University.

that paper in two important cases: (i) when the action is isometric and (ii) when the compact quantum group is finite dimensional. We also mention the work of Etingof and Walton [4] which gives a similar no go result in the algebraic framework.

It is, however, expected that genuine non-compact quantum groups may have faithful smooth actions on smooth connected manifolds. Indeed, there are examples of such actions in the algebraic set-up (see [6, Example 14.2]). This motivates one to construct examples of faithful C*-actions of non-compact locally compact quantum groups on $\mathrm{C}_0(X)$ where X is a smooth connected manifold. It is also desirable to see if the actions are smooth in any suitable sense. We give a method to construct such actions using the theory of bicrossed products due to Baaj, Skandalis and Vaes [1] and Vaes and Vainermann [15]. There is one remarkable observation: there are several non-Kac quantum groups with faithful C*-actions (even ergodic) on $\mathrm{C}_0(X)$. This could not be possible in the realm of compact quantum groups, as any compact quantum group acting faithfully on a commutative C*- algebra must be of Kac type (see [8]). Thus, there seems to be more freedom for getting quantum symmetries on classical spaces in the realm of locally compact quantum groups than the compact ones. On the other hand, it should be noted that if we are interested in actions which are isometric in a natural sense (as in [5]). We cannot possibly get any genuine locally compact quantum group actions on classical (connected) Riemannian manifolds. Indeed, such a no-go result is obtained within the class of locally compact quantum groups considered by us in this paper (see Theorem 4.7).

The plan of the paper is as follows. We gather some basic definitions and facts about locally compact quantum groups and their actions in the von Neumann as well as C*-algebraic set-up in Subsection 2.1, followed by a brief account of the bicrossed product construction of locally compact groups in Subsection 2.2. In Section 3, we specialise to the locally compact quantum groups arising from bicrossed product construction of two groups say G_1, G_2 forming a matched pair, with G_1 being an abelian Lie group. We describe a natural action of this bicrossed product quantum group on $\mathrm{C}_0(\widehat{G_1})$ and verify that it satisfies Podleś-type density conditions. Section 4 is devoted to investigate necessary and sufficient conditions for this action to be faithful or isometric. Using this, we observe that that there is a large class of genuine locally compact quantum groups having faithful actions on commutative C*-algebra of C_0-functions on a locally compact manifold in Section 5. However, it is shown in Subsection 4.2 that no such genuine quantum group actions can be isometric.

2. Preliminaries

All Hilbert spaces and C*-algebras are assumed to be separable.

For two norm-closed subsets X and Y of a C*-algebra, let

$$X \cdot Y := \{xy : x \in X, y \in Y\}^{\mathrm{CLS}},$$

where CLS stands for the *closed linear span*.

For a C*-algebra A, let $\mathcal{M}(A)$ be its multiplier algebra and $\mathcal{U}(A)$ be the group of unitary multipliers of A. The unit of $\mathcal{M}(A)$ is denoted by 1_A. Next recall some standard facts about multipliers and morphisms of C*-algebras from [13, Appendix A]. Let A and B be C*-algebras. A *-homomorphism $\varphi\colon A \to \mathcal{M}(B)$ is called *nondegenerate* if $\varphi(A) \cdot B = B$. Each nondegenerate *-homomorphism $\varphi\colon A \to \mathcal{M}(B)$ extends uniquely to a unital *-homomorphism $\widetilde{\varphi}$ from $\mathcal{M}(A)$ to $\mathcal{M}(B)$. Let $\mathfrak{C}^*\mathfrak{alg}$ be the category of C*-algebras with nondegenerate *-homomorphisms $A \to \mathcal{M}(B)$ as morphisms $A \to B$; let Mor(A,B) denote this set of morphisms. We use the same symbol for an element of $\mathrm{Mor}(A, B)$ and its unique extension from $\mathcal{M}(A)$ to $\mathcal{M}(B)$.

A *representation* of a C*-algebra A on a Hilbert space $\mathcal{H}$ is a nondegenerate *-homomorphism $\pi\colon A \to \mathbb{B}(\mathcal{H})$. Since $\mathbb{B}(\mathcal{H}) = \mathcal{M}(\mathbb{K}(\mathcal{H}))$, the nondegeneracy conditions $\pi(A)\cdot\mathbb{K}(\mathcal{H}) = \mathbb{K}(\mathcal{H})$ is equivalent to begin $\pi(A)(\mathcal{H})$ is norm dense in $\mathcal{H}$, and hence this is same as having a morphism from A to $\mathbb{K}(\mathcal{H})$. The identity representation of $\mathbb{K}(\mathcal{H})$ on $\mathcal{H}$ is denoted by $\mathrm{id}_{\mathcal{H}}$. The group of unitary operators on a Hilbert space $\mathcal{H}$ is denoted by $\mathcal{U}(\mathcal{H})$. The identity element in $\mathcal{U}(\mathcal{H})$ is denoted by $1_{\mathcal{H}}$.

We use $\otimes$ both for the tensor product of Hilbert spaces, minimal tensor product of C*-algebras, and von Neumann algebras which is well understood from the context. We write Σ for the tensor flip $\mathcal{H}\otimes\mathcal{K}\to\mathcal{K}\otimes\mathcal{H}$, $x\otimes y\mapsto y\otimes x$, for two Hilbert spaces $\mathcal{H}$ and $\mathcal{K}$. We write σ for the tensor flip isomorphism $A\otimes B\to B\otimes A$ for two C*-algebras or von Neumann algebras A and B.

Let A_1, A_2, A_3 be C*-algebras. For any $t\in\mathcal{M}(A_1\otimes A_2)$ we denote the leg numberings on the level of C*-algebras as $t_{12} := t\otimes 1_{A_3}\in\mathcal{M}(A_1\otimes A_2\otimes A_3)$, $t_{23} := 1_{A_3}\otimes t_{12}\in\mathcal{M}(A_3\otimes A_1\otimes A_2)$ and $t_{13} := \sigma_{12}(t_{23}) = \sigma_{23}(t_{12})\in\mathcal{M}(A_1\otimes A_3\otimes A_2)$. In particular, let $A_i = \mathbb{B}(\mathcal{H}_i)$ for some Hilbert spaces $\mathcal{H}_i$, where $i = 1,2,3$. Then for any $t\in\mathbb{B}(\mathcal{H}_1\otimes\mathcal{H}_2)$ the leg numberings are obtained by replacing σ with the conjugation by Σ operator.

2.1. **Locally compact quantum groups and their actions.** For a general theory of C*-algebraic locally compact quantum groups we refer [11, 13].

A C*-*bialgebra* (A,Δ_A) is a C*-algebra A and a comultiplication $\Delta_A\in\mathrm{Mor}(A, A\otimes A)$ that is coassociative: $(\mathrm{id}_A\otimes\Delta_A)\circ\Delta_A = (\Delta_A\otimes\mathrm{id}_A)\circ\Delta_A$. Moreover, (A,Δ_A) is *bisimplifiable* C*-bialgebra if Δ_A satisfies the cancellation property

$$\Delta_A(A)\cdot(1_A\otimes A) = \Delta_A(A)\cdot(A\otimes 1_A) = A\otimes A. \tag{2.1}$$

Let φ be a faithful (approximate) KMS weight on A (see [10, Section 1]). The set of all positive φ-integrable and φ-square integrable elements are defined by $\mathcal{M}_\varphi^+ := \{a\in A^+ \mid \varphi(a)<\infty\}$ and $\mathcal{N}_\varphi := \{a\in A\mid \varphi(a^*a)<\infty\}$, respectively. Moreover, φ is called

(1) *left invariant* if $\omega((\mathrm{id}_A\otimes\varphi)\Delta_A(a)) = \omega(1)\varphi(a)$ for all $\omega\in A_*^+$, $a\in\mathcal{M}_\varphi^+$;
(2) *right invariant* if $\omega((\varphi\otimes\mathrm{id}_A)\Delta_A(a)) = \omega(1)\varphi(a)$ for all $\omega\in A_*^+$, $a\in\mathcal{M}_\varphi^+$.

Definition 2.2 ([10, Definition 4.1])**.** A *locally compact quantum group* (*quantum groups* from now onwards) is a bisimplifiable C*-bialgebra $\mathbb{G} = (A,\Delta_A)$ with left and right invariant approximate KMS weights φ and ψ, respectively.

By [10, Theorem 7.14 & 7.15], invariant weights φ and ψ are unique up to a positive scalar factor; hence they are called the left and right *Haar weights* for $\mathbb{G}$. Moreover, there is a unique (up to isomorphism) Pontrjagin dual $\widehat{\mathbb{G}} = (\hat{A},\hat{\Delta}_A)$ of $\mathbb{G}$, which is again a quantum group.

Next we consider the GNS triple $(\mathrm{L}^2(\mathbb{G}),\pi,\Lambda)$ for φ. There is an element $\mathbb{W}\in\mathcal{U}(\mathrm{L}^2(\mathbb{G})\otimes\mathrm{L}^2(\mathbb{G}))$ satisfying the pentagon equation:

$$\mathbb{W}_{23}\mathbb{W}_{12} = \mathbb{W}_{12}\mathbb{W}_{13}\mathbb{W}_{23} \qquad \text{in } \mathcal{U}(\mathrm{L}^2(\mathbb{G})\otimes\mathrm{L}^2(\mathbb{G})\otimes\mathrm{L}^2(\mathbb{G})). \tag{2.3}$$

$\mathbb{W}$ is called *multiplicative unitary*. Furthermore, [10, Proposition 6.10] shows that $\mathbb{W}$ is manageable (in the sense of [18, Definition 1.2]). Also $\mathbb{W}$ *generates* $\mathbb{G}$ (or $\mathbb{G}$ is *generated* by $\mathbb{W}$) in the following sense:

(1) the dual multiplicative unitary $\widehat{\mathbb{W}} := \Sigma\mathbb{W}^*\Sigma\in\mathcal{U}(\mathrm{L}^2(\mathbb{G})\otimes\mathrm{L}^2(\mathbb{G}))$ is also manageable.
(2) the slices of $\mathbb{W}$ defined by

$$\hat{A} := \{(\omega\otimes\mathrm{id}_{\mathrm{L}^2(\mathbb{G})})\mathbb{W} : \omega\in\mathbb{B}(\mathrm{L}^2(\mathbb{G}))_*\}^{\mathrm{CLS}}, \tag{2.4}$$

$$A := \{(\mathrm{id}_{\mathrm{L}^2(\mathbb{G})}\otimes\omega)\mathbb{W} : \omega\in\mathbb{B}(\mathrm{L}^2(\mathbb{G}))_*\}^{\mathrm{CLS}}, \tag{2.5}$$

are nondegenerate C*-subalgebras of $\mathbb{B}(\mathrm{L}^2(\mathbb{G}))$.

(3) $\mathbb{W} \in \mathcal{U}(A \otimes \hat{A}) \subseteq \mathcal{U}(\mathrm{L}^2(\mathbb{G}) \otimes \mathrm{L}^2(\mathbb{G}))$.

(4) the comultiplication maps Δ_A and $\hat{\Delta}_A$ are defined by

$$\Delta_A(a) := \mathbb{W}^*(1 \otimes a)\mathbb{W}, \qquad \hat{\Delta}_A(\hat{a}) := \sigma\big(\mathbb{W}(\hat{a} \otimes 1)\mathbb{W}^*\big), \tag{2.6}$$

for all $a \in A$, $\hat{a} \in \hat{A}$.

A general theory of locally compact quantum groups in the von-Neumann algebraic framework has been developed by Kustermans and Vaes in [11]. Moreover, there is a nice interplay between C*-algebraic and von-Neumann algebraic locally compact quantum groups in general via multiplicative unitaries. We briefly recall this in the group case.

Let G be a locally compact group and μ be its left Haar measure. Define, $\mathbb{W} \in \mathcal{U}(L^2(G \times G, \mu \times \mu)$ by $\mathbb{W}\xi(x, y) = \xi(x, x^{-1}y)$ for all $\xi \in L^2(G \times G, \mu \times \mu)$, $x, y \in G$. A simple computation shows that $\mathbb{W}$ is a multiplicative unitary. Furthermore,

$$A = \mathrm{C}_0(G) = \{(\mathrm{id}_{\mathrm{L}^2(G,\mu)} \otimes \omega)\mathbb{W} : \omega \in \mathbb{B}(\mathrm{L}^2(G))_*\}^{\mathrm{CLS}},$$
$$M = L^\infty(G) = \{(\mathrm{id}_{\mathrm{L}^2(\mathbb{G})} \otimes \omega)\mathbb{W} : \omega \in \mathbb{B}(\mathrm{L}^2(\mathbb{G}))_*\}^{\text{weak closure}}.$$

Using (2.6) we get $\Delta_M \colon L^\infty(G) \to L^\infty(G \times G)$ by $\Delta_M(f)(x, y) = f(xy)$ for all $f \in L^\infty(G)$. The pair (M, Δ_M) is a von-Neumann bialgebra. The left Haar weight φ on $L^\infty(G)$ is given by the integration with respect to the left Haar measure μ on G: $\varphi(f) := \int f(h)\mathrm{d}\mu(h)$ for $f \in L^\infty(G_2)^+$. Similarly, the right Haar weight ψ on (M, Δ_M) is obtained from the to the right Haar measure on G. The pair $\mathbb{G} = (M, \Delta_M)$ is the von Neumann algebraic locally compact quantum group associated to G.

Similarly, $\Delta_A \colon \mathrm{C}_0(G) \to \mathrm{C}_\mathrm{b}(G \times G)$ is obtained by restricting Δ_M on $\mathrm{C}_0(G)$. The restriction of left and right Haar weight of (M, Δ_M) on $\mathrm{C}_0(G)$ defines the left and Haar weight on (A, Δ_A), respectively. Thus (A, Δ_A) is the C*-algebraic version of $\mathbb{G}$.

Let $\hat{A} = \mathrm{C}^*_\mathrm{r}(G)$ and $\hat{M} = L(G)$. Let λ be the left regular representation of G on $L^2(G, \mu)$. Define, $\hat{\Delta}_M \colon L(G) \to L(G \times G)$ by $\hat{\Delta}(\lambda_g) = \lambda_g \otimes \lambda_g$ and $\hat{\Delta}_A \colon \mathrm{C}^*_\mathrm{r}(G) \to \mathcal{M}(\mathrm{C}^*_\mathrm{r}(G \times G))$ as the restriction of $\hat{\Delta}_M$. The left and right invariant invariant Haar weights on $\hat{M}$ is given by $\hat{\varphi}(\lambda(f)) = f(1_G)$ for all $f \in \mathrm{C}_\mathrm{c}(G)$ such that $\lambda(f) \in L(G)^+$. Then $\widehat{\mathbb{G}} = (\hat{A}, \hat{\Delta}_A)$ and $\widehat{\mathbb{G}} = (\hat{M}, \hat{\Delta}_M)$ are the dual of $\mathbb{G}$ in C*-algebraic and von Neumann algebraic setting, respectively.

Definition 2.7. A (*right*) C*-*action* of $\mathbb{G}$ on a C*-algebra C is a morphism $\gamma \colon C \to C \otimes A$ with the following properties:

(1) γ is a comodule structure, that is,

$$(\mathrm{id}_C \otimes \Delta_A)\gamma = (\gamma \otimes \mathrm{id}_A)\gamma; \tag{2.8}$$

(2) γ satisfies the *Podleś condition*:

$$\gamma(C) \cdot (1_C \otimes A) = C \otimes A. \tag{2.9}$$

Some authors demands γ to be injective. Similarly, the von Neumann algebraic version (M, Δ_M) of $\mathbb{G}$ acts on von Neumann algebras as well.

Definition 2.10. A (*right*) *von Neumann algebraic action* of $\mathbb{G}$ on a von Neumann algebra N is a faithful, normal, unital *-homomorphism $\gamma \colon N \to N \otimes M$ satisfying $(\mathrm{id}_N \otimes \Delta_M)\gamma = (\gamma \otimes \mathrm{id}_M)\gamma$.

2.2. **Bicrossed product of groups.** Bicrossed product construction for mathced pair of locally compact quantum groups goes back to the work of Baaj and Vaes [2] and Vaes and Vainerman [15]. In this article, we shall restrict out attention to the bicrossed product construction for locally compact groups.

Definition 2.11 ([15, Definition 4.7])**.** Let G_1, G_2 and G be locally compact groups with fixed left Haar measures. The pair (G_1, G_2) is called a *matched pair* if

(1) there exist a homomorphism $i\colon G_1 \hookrightarrow G$ and an anti-homomorphism $j\colon G_2 \hookrightarrow G$ with closed images and homeomorphism onto these images;
(2) $\theta\colon G_1 \times G_2 \to G$ defined by $\theta((g,h)) = i(g)j(h)$ is an homeomorphism onto an open subgroup Ω of G having a complement of measure zero.

This allows to define almost everywhere and measurable left action $(\alpha_g)_{g\in G_1}$ of G_1 on G_2 and right action $(\beta_h)_{h\in G_2}$ on G, and satisfies $j(\alpha_g(h))i(\beta_h(g)) = i(g)j(h)$ for almost all $g \in G_1$ and $h \in G_2$.

By [15, Lemma 4.9], the maps $G_1 \times G_2 \to G_2\colon (g,h) \mapsto \alpha_g(h)$ and $G_1 \times G_2 \to G_1\colon (g,h) \mapsto \beta_h(g)$ are measurable, defined almost everywhere, and satisfy the following relations:

$$\alpha_{gs}(h) = \alpha_g(\alpha_s(h)), \qquad \beta_h(gs) = \beta_{\alpha_s(h)}(g)\beta_h(s), \tag{2.12}$$

$$\beta_{ht}(g) = \beta_h(\beta_t(g)), \qquad \alpha_g(ht) = \alpha_{\beta_t(g)}(h)\alpha_g(t). \tag{2.13}$$

for almost all $g, s \in G_1$, $h, t \in G_2$. Also, $\alpha_g(1_{G_2}) = 1_{G_2}$, and $\beta_h(1_{G_1}) = 1_{G_1}$ for all $g \in G$, $h \in H$.

Then $\alpha\colon L^\infty(G_2) \to L^\infty(G_1 \times G_2)$ defined by $\alpha(f)(g,s) := f(\alpha_g(s))$ is a (left) von Neumann algebraic action of the locally compact quantum group $\mathbb{G}_1 = ((L^\infty(G_1), \Delta_{L^\infty(G_1)})$ on $L^\infty(G_2)$. The von Neumann algebraic version of the bicrossed product $\mathbb{G} = (M, \Delta_M)$ is given by

$$M := \big(\alpha(L^\infty(G_2))(L(G_1) \otimes 1)\big)'', \qquad \Delta_M(z) := \mathbb{W}^*(1 \otimes z)\mathbb{W} \tag{2.14}$$

Let λ be the left regular representation of G_1. The left (and right) Haar weight on $L^\infty(G_2)$. Let $\hat{\alpha}$ be the dual action of $L(G_1)$ on the crossed product M. Then $\hat{\alpha}\colon M \to L(G_1) \otimes M$ is defined by $\hat{\alpha}(\alpha(y)) = 1_{L(G_1)} \otimes \alpha(\eta)$ and $\hat{\alpha}(x \otimes 1_{L^\infty(G_2)}) = \hat{\Delta}_{L(G_1)}(x) \otimes 1_{L^\infty(G_2)}$ for all $x \in L(G_1)$ and $y \in L^\infty(G_2)$.

Let $\hat{\varphi}_1$ and φ_2 be the left Haar weights on $L(G_1)$ and $L^\infty(G_2)$, respectively. By [15, Definition 1.13], the left Haar weight φ on $\mathbb{G}$ is given by $\varphi := \varphi_2\alpha^{-1}(\hat{\varphi}_1 \otimes \mathrm{id} \otimes \mathrm{id})\hat{\alpha}$. A simple computation shows, for any $f \in \mathcal{N}_{\hat{\varphi}_1}$ and $\eta \in \mathcal{N}_{\varphi_2}$,

$$\varphi\big((\alpha(\eta)(\lambda(f) \otimes 1)\big) = \hat{\varphi}_1(\lambda(f))\varphi_2(\eta) = \varphi\big((\lambda(f) \otimes 1)(\alpha(\eta)\big) \tag{2.15}$$

Let $\mathcal{H} = L^2(G_1 \times G_2)$ be the Hilbert space of square integrable functions with respect to the product of the left Haar measures of G_1 and G_2. Using [1, Definition 3.3] for left Haar measure, we obtain a multiplicative unitary $\mathbb{W} \in \mathcal{U}(\mathcal{H} \otimes \mathcal{H})$ for $\mathbb{G}$ defined by

$$\mathbb{W}\xi(g,s,h,t) := \xi(\beta_{\alpha_g(s)^{-1}t}(h)g, s, h, \alpha_g(s)^{-1}t), \tag{2.16}$$

for $\xi \in L^2(G_1 \times G_2 \times G_1 \times G_2)$, and for almost all $g, s \in G_1$, $h, t \in G_2$.

Finally, we recall the C*-algebraic version of $\mathbb{G}$ from [1, Section 3]. Equip the quotient space $G_1\backslash G$ with its canonical invariant measure class. Then embedding $G_2 \to G_1\backslash G$ identifies G_2 with a Borel subset of $G_1\backslash G$ with complement of measure zero. Then [1, Proposition 3.2] gives an isomorphism between $L^\infty(G_2)$ and $L^\infty(G_1\backslash G)$; hence we can restrict α to $\mathrm{C}_0(G_1\backslash G)$. The C*-algebraic version (A, Δ_A) of $\mathbb{G}$ is given by [1, Proposition 3.6]

$$A := \alpha(\mathrm{C}_0(G_1\backslash G)) \cdot (\lambda(\mathrm{C}^*_{\mathrm{r}}(G_1)) \otimes 1), \qquad \alpha \in \mathrm{Mor}(\mathrm{C}_0(G_2), A). \tag{2.17}$$

The dual $\widehat{\mathbb{G}} = (\hat{A}, \hat{\Delta}_A)$ is obtained by exchanging the roles of G_1 and α by G_2 and β, respectively.

3. Existence of C*-actions of bicrossed products on spaces

Let G_1, G_2 be locally compact groups and (G_1, G_2) form a matched pairs. Let $\mathbb{G} = (A, \Delta_A)$ be the associated (C*-algebraic) bicrossed product quantum group. From now onwards we assume that G_1 is abelian.

Theorem 3.1. *There is a* C*-*action of* $\mathbb{G}$ *on* $\mathrm{C}_0(\widehat{G_1})$.

Throughout λ denotes the (right) regular representation of G_1: $\lambda(f)\xi'(h) := \int f(g)\xi'(hg)\mathrm{d}\mu_1(g)$, for $f \in \mathrm{C_c}(G_1)$ and $\xi' \in L^2(G_1)$.

Clearly, we have an element $i \in \mathrm{Mor}(\mathrm{C}^*_{\mathrm{r}}(G_1), A)$ and its extension, denoted by i again, to $\mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1))$ is given by $i(x) := x \otimes 1$ for all $x \in \mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1))$. Hence, $\gamma := \Delta_A \circ i$ is an element in $\mathrm{Mor}(\mathrm{C}^*_{\mathrm{r}}(G_1), A \otimes A)$. In order to interpret γ as the desired C*-action in Theorem 3.1 following proposition will be crucial.

Proposition 3.2. *γ is a *-homomorphism from $\mathrm{C}^*_{\mathrm{r}}(G_1)$ to $\mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1) \otimes A)$ and defined by*

$$\gamma(\lambda(f))\xi(g,h,t) := \int f(z)\xi(g\beta_{\alpha_h(t)}(z), hz, t)\mathrm{d}\mu_1(z) \tag{3.3}$$

for all $f \in \mathrm{C_c}(G_1)$ and $\xi \in L^2(G_1 \times G_1 \times G_2)$.

Proof. The adjoint $\mathbb{W}^* \in \mathcal{U}(L^2(G_1 \times G_2 \times G_1 \times G_2))$ of the multiplicative unitary $\mathbb{W}$ of $\mathbb{G}$ in (2.16) is defined by

$$\mathbb{W}^*\xi'(g,s,h,t) := \xi'(g', s, h, \alpha_{g'}(s)t) \qquad \text{for } \xi' \in L^2(G_1 \times G_2 \times G_1 \times G_2),$$

where $g' := \beta_t(h)^{-1}g$ for all $g, h \in G_1$ and $s, t \in G_2$. Using the definition of the comultiplication 2.6 and the property (2.13) of β we obtain

$$\begin{aligned}
\gamma(\lambda(f))\xi'(g,s,h,t) &= \Delta_A(1_{L^2(G_1\times G_2)} \otimes \lambda(f) \otimes 1_{L^2(G_2)})\xi'(g,s,h,t)\\
&= \big(\mathbb{W}^*(1_{L^2(G_1\times G_2)} \otimes \lambda(f) \otimes 1_{L^2(G_2)})\mathbb{W}\big)\xi'(g,s,h,t)\\
&= \big((1_{L^2(G_1\times G_2)} \otimes \lambda(f) \otimes 1_{L^2(G_2)})\mathbb{W}\big)\xi'(g',s,h,\alpha_{g'}(s)t)\\
&= \mathbb{W}\Big(\int f(z)\xi'(g',s,hz,\alpha_{g'}(s)t)\mathrm{d}\mu_1(z)\Big)\\
&= \int f(z)\xi'(g\beta_{\alpha_h(t)}(z), s, hz, t)\mathrm{d}\mu_1(z).
\end{aligned} \tag{3.4}$$

Since G_1 is abelian, we indenify $\mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1) \otimes A)$ with $\mathrm{C_b}(\widehat{G_1}, \mathcal{M}(A)) \subset \mathbb{B}(L^2(\widehat{G_1} \times G_1 \times G_2))$ using the Fourier transform.

Let $f\colon \widehat{G_1} \to \mathcal{M}(A)$ be a strictly continuous function. Define an operator M_f acting on $L^2(\widehat{G_1} \times G_1 \times G_2)$ by

$$M_f(\xi_1 \otimes \xi_2)(\hat{g}, h, t) := \xi_1(\hat{g})(f(\hat{g})\xi_2)(h,t) \quad \text{for all } \xi_1 \in L^2(\widehat{G_1}),\ \xi_2 \in L^2(G_1 \times G_2),$$

which is an element in $M_f \in \mathrm{C_b}(\widehat{G_1}, \mathcal{M}(A)) \cong \mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1) \otimes A)$.

Recall the dual pairing $\langle \cdot, \cdot \rangle\colon G_1 \times \widehat{G_1} \to \mathbb{T}$ defined by $\langle h, \hat{g} \rangle := \hat{g}(h)$ for all $h \in G_1$ and $\hat{g} \in \widehat{G_1}$. Let D be a C*-algebra. For an element $F \in L^1(G_1, D)$, the Fourier transform $\widehat{F}$ is defined by $\widehat{F}(\hat{g}) := \int F(h)\langle h, \hat{g}\rangle \mathrm{d}\mu_1(h)$ for $\hat{g} \in \widehat{G_1}$, and $\widehat{F}$ is an element in $\mathrm{C}_0(\widehat{G_1}, D)$.

Then, for a fixed $z \in G_1$, consider $T_z\colon \widehat{G_1} \to \mathcal{M}(A)$ defined by

$$T_z(\hat{g})\xi_2(h,t) := \langle \beta_{\alpha_h(t)}(z), \hat{g}\rangle \xi_2(hz, t) \qquad \text{for all } \xi_2 \in L^2(G_1 \times G_2).$$

Clearly, $\hat{g} \mapsto T_z(\hat{g})$ is continuous in the strict topology. Hence, any $f \in \mathrm{C_c}(G_1)$ gives $\int f(z)T_z\mathrm{d}\mu_1(z)$ is an element in $\mathrm{C_b}(\widehat{G_1}, \mathcal{M}(A)) \cong \mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1) \otimes A)$.

Using the Fourier transform in the first leg of (3.4) we observe

$$\gamma(\lambda(f)) = \Sigma_{12}(\int f(z)(\mathrm{id}_{L^2(G_2)} \otimes T_z \mathrm{d}\mu_1(z))\Sigma_{12}$$

in $\mathbb{B}(L^2(\widehat{G_1} \times G_2 \times G_1 \times G_2))$. Finally, putting $s = 1_{G_2}$ in (3.4) we obtain (3.3). □

We gather some standard facts related to Fourier transform in the next lemma.

Lemma 3.5. *Let K be a compact subset of G_1. Define*

$$S(K, D) := \{F \in L^1(G_1, D) \mid \mathrm{supp}(F) \subset K\} \subset L^1(G, D);$$
$$\widehat{S(K, D)} := \{\widehat{F} \mid F \in S(K, D)\} \subset \mathrm{C}_0(\widehat{G_1}, D).$$

We have the following:

1. *Let $\widehat{G_1}$ is a Lie group, and let $\delta = (\delta_1, \cdots, \delta_n)$ be the canonical derivation on $\widehat{G_1}$ for $i = 1, \cdots, n$. Then $\widehat{S(K, D)} \subset \mathrm{Dom}(\tilde{\delta})$ and $\tilde{\delta}\big(\widehat{S(K, D)}\big) \subset \widehat{S(K, D)}$, where $\tilde{\delta}_i := \delta_i \otimes \mathrm{id}_D$ for $i = 1, \cdots, n$, and $\tilde{\delta} := (\tilde{\delta}_1, \cdots, \tilde{\delta}_n)$.*
2. *$S(K, D), \widehat{S(K, D)} \subset L^2(G_1, D)$.*
3. *Let D' be a C^*-algebra and $\rho\colon D \to D'$ be a completely bounded map. Then $(\mathrm{id} \otimes \rho)S(K, D) \subset S(K, D')$ and $(\mathrm{id} \otimes \rho)\widehat{S(K, D)} \subset \widehat{S(K, D')}$.*

Proof. By definition $\delta_i(\lambda_g) := \rho_i(g)\lambda_g$, where $\lambda_g \in \mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1)) \cong \mathrm{C}_{\mathrm{b}}(\widehat{G_1})$ and $\rho_i \in \widehat{G_1}$ for $i = 1, \cdots, n$; hence gives (1). The second fact follows because the Fourier transform is L^2-isometry. The last fact is trivial. □

For any $f \in \mathrm{C}_{\mathrm{c}}(G_1)$ and $\eta \in \mathrm{C}_{\mathrm{c}}(G_2)$ define $\hat{\pi}_1(f) := \lambda(f) \otimes 1$ and $\pi_2(\eta) := \alpha(\eta)$. Therefore, $\hat{\pi}_1(f)\pi_2(\eta) \in A$.

Proposition 3.6. *Define $w := \gamma(\hat{\pi}_1(f))(1 \otimes \pi_2(\eta)) \in S(K, \mathcal{M}(A))$, where $K = \mathrm{supp}(f)$. Then $p \mapsto (\mathrm{id} \otimes \varphi)(\hat{w}(p)^*\hat{w}(p)) \in \mathrm{C}_0(\widehat{G_1})$.*

Proof. By definition

$$w^*w = (1 \otimes \pi_2(\eta^*))\gamma\big(\hat{\pi}_1(|f|^2)\big)(1 \otimes \pi_2(\eta))$$

Using (2.15) we get

$$\begin{aligned}(\mathrm{id} \otimes \varphi)(w^*w) &= (\mathrm{id} \otimes \varphi)\big(1 \otimes \pi_2(\eta^*))\gamma(\hat{\pi}_1(|f|^2))(1 \otimes \pi_2(\eta)\big) \\ &= (\mathrm{id} \otimes \varphi)\big(1 \otimes \pi_2(|\eta|^2)\gamma(\hat{\pi}_1(|f|^2)\big)\end{aligned}$$

By virtue of Proposition 3.2 and (2.15), for $p \in \widehat{G_1}$, we get

$$\begin{aligned}(\mathrm{id} \otimes \varphi)(\hat{w}(p)^*\hat{w}(p)) &= \iint |f|^2(g)|\eta|^2(s)\langle \beta_s(g), p\rangle \mathrm{d}\mu_1(g)\mathrm{d}\mu_2(s) \\ &= \iint |f|^2(\beta_{s^{-1}}(g'))|\eta|^2(s)\theta(g', s)\langle g', p\rangle \mathrm{d}\mu_1(g')\mathrm{d}\mu_2(s),\end{aligned}$$

where $g' = \beta_s(g)$ and $\theta(g, s) := |\frac{\mathrm{d}}{\mathrm{d}\mu_1}\beta_{s^{-1}}(g)|$.

Define $G_s(g) := |f|^2(\beta_{s^{-1}}(g))|\eta|^2(s)\theta(g, s)$. Then,

$$(\mathrm{id} \otimes \varphi)(\hat{w}(p)^*\hat{w}(p)) = \int \widehat{G_s}(p)\mathrm{d}\mu_2(s).$$

A simple computation gives

$$\begin{aligned}\|G_s\|_1 = \int \|G_s(g)\|\mathrm{d}\mu_1(g) &= \int |f|^2(\beta_{s^{-1}}(g))|\eta|^2(s)\theta(g, s)\mathrm{d}\mu_1(g) \\ &= \int |f|^2(g)|\eta|^2(s)\mathrm{d}\mu_1(g) = (\|f\|_2)^2|\eta|^2(s).\end{aligned}$$

Therefore, $\widehat{G_s} \in \mathrm{C}_0(\widehat{G_1})$ for almost all $s \in G_2$. Also $\int |\eta|^2(s)\mathrm{d}\mu_2(s) \leq \infty$. By dominated convergence theorem, for any sequence $\{p_n\} \subset \widehat{G_1}$ such that $|p_n| \to \infty$ as $n \to \infty$, we have

$$\lim_{n\to\infty} \int \widehat{G_s}(p_n)\mathrm{d}\mu_2(s) = \int \big(\lim_{n\to\infty} \widehat{G_s}(p_n)\big)\mathrm{d}\mu_2(s) = 0. \qquad \square$$

Proof of Theorem 3.1. Proposition 3.2 and Proposition 3.6 give γ is an element in $\mathrm{Mor}(\mathrm{C}_0(\widehat{G_1}), \mathrm{C}_0(\widehat{G_1}) \otimes A)$. The coassociativity of Δ_A gives (2.8) for γ.

Let $(L^2(\mathbb{G}), \pi, \Lambda)$ be the GNS triple for the the left Haar weight φ in (2.15). For any $v \in L^2(\mathbb{G})$, define the operator $\Theta_v(c) := cv$ for $c \in \mathbb{C}$. Let $(e_i)_{i\in\mathbb{N}}$ be an orthonormal basis of $L^2(\mathbb{G})$. For $w \in S(K, \mathcal{M}(A))$ in Proposition 3.6, define

$$x_i^*(p) := (\mathrm{id} \otimes \Theta_{e_i}^*)(\mathrm{id} \otimes \Lambda)\Delta_A(\hat{w}(p)) \in \mathcal{M}(A) \qquad \text{for } p \in \widehat{G_1}.$$

Also, for $q \in \mathcal{N}_\varphi$ define $q_i := (\mathrm{id} \otimes \Theta_{e_i}^*)(\mathrm{id} \otimes \Lambda)\Delta_A(q) \in \mathcal{M}(A)$.

We compute,

$$\begin{aligned}\sum_{i=1}^{\infty} x_i(p)q_i &= \sum_{i=1}^{\infty} (\mathrm{id} \otimes \Lambda)\Delta_A(w(p))^*(1 \otimes \Theta_{e_i}\Theta_{e_i}^*)(\mathrm{id} \otimes \Lambda)\Delta_A(q) \\ &= (\mathrm{id} \otimes \Lambda)\Delta_A(w(p))^*(\mathrm{id} \otimes \Lambda)\Delta_A(q) \\ &= \varphi(\hat{w}(p)^*q)1_{\mathcal{M}(A)}\end{aligned}$$

By virtue of Proposition 3.6, $F_N(p) := \Sigma_{i=1}^N x_i(p)q_i$ is strictly convergent. Hence, for any given $q' \in A$, the sequence $\{F_N(1 \otimes q')\}$ converges uniformly over every compact subset of $\widehat{G_1}$.

In order to establish Podleś condition (2.9) for γ we need to show $\{F_N(1 \otimes q')\}$ converges uniformly over $\widehat{G_1}$ for $q' \in A$.

By a similar argument used in [16, Proposition 5.11], and Proposition 3.6 gives $\sum_{i=1}^n x_i^*(p)x_i(p)$ strictly converges to $\varphi(\hat{w}(p)^*\hat{w}(p))1_{\mathcal{M}(A)}$. Similarly, $\sum_{i=1}^n q_i^*q_i$ is bounded and strictly convergent. Let $\|\sum_{i=1}^n q_i^*q_i\| < C^2$. Given $\epsilon > 0$, we can choose a compact subset K' of $\widehat{G_1}$, such that $(\mathrm{id} \otimes \varphi)(\hat{w}^*(p)\hat{w}(p)) \leq (\frac{\epsilon}{C})^2$ for all $p \notin K'$. Hence $\|\sum_{i=m}^n x_i(p)x_i^*(p)\| \leq (\frac{\epsilon}{C})^2$ for all $p \notin K'$, and for all m, n.

Now choose N_0 such that for all $m, n \geq N_0$, $\|(F_m - F_n)(p)q'\| < \epsilon$ for $p \in K'$, $q' \in A$.

Finally, for all $m, n \geq N_0$

$$\|(F_m - F_n)(p)\| \leq \left\|\sum_{i=m}^n x_i(p)x_i^*(p)\right\|^{\frac{1}{2}} \left\|\sum_{i=m}^n q_i^*q_i\right\|^{\frac{1}{2}} < \epsilon\|q'\|,$$

for $p \notin K'$. Hence $\{F_N(1 \otimes q')\}$ is Cauchy sequence in norm for $q' \in A$. $\square$

4. Properties of bicrossed product C*-actions

Let G_1, G_2, and $\mathbb{G} = (A, \Delta_A)$ be as before, so that G_1 is abelian. In this section we shall discuss various properties of the C*-action γ of $\mathbb{G}$ on $\mathrm{C}_0(\widehat{G_1})$ constructed in Theorem 3.1.

Recall, a right action γ of a von Neumann algebraic quantum group $\mathbb{G} = (M, \Delta_M)$ on a von Neumann algebra N is called *ergodic* if $N^\gamma := \{x \in N : \ \gamma(x) = x \otimes 1_M\}$ is equal to $\mathbb{C} \cdot 1_N$.

Proposition 4.1. *The von Neumann algebraic action γ of $\mathbb{G}$ on $L(G_1)$ is ergodic.*

Proof. Let (M, Δ_M) be the von Neumann algebraic version of $\mathbb{G}$. By construction, γ is obtained from the comultiplication Δ_A of the C*-algebraic version of $\mathbb{G}$. Since, Δ_A extends uniquely to the comultiplication Δ_M on M by (2.14). Then, in a similar way, γ also extends to a von Neumann algebraic action, denoted again by γ, of $\mathbb{G}$

on $L(G_1)$. Now ergodicity of Δ_M (see [10, Result 5.13] or [14, Theorem 2.1]) implies the same for γ. □

4.1. **Faithfulness.** Motivated by [6, Definition 4.5] we propose the following definition, as a possible generalisation of faithful actions on locally compact quantum groups on C*-algebras.

Definition 4.2. A C*-action $\gamma\colon C \to C \otimes A$ of $\mathbb{G} = (A, \Delta_A)$ on a C*-algebra C is called *faithful* if the *-algebra generated by $\{(\omega \otimes \mathrm{id}_A)\gamma(c) : \omega \in C',\, c \in C\}$ is strictly dense in $\mathcal{M}(A)$.

Example 4.3. In particular, the comultiplication map Δ_A is a C*-action of $\mathbb{G}$ on A. Given any $\omega \in A'$ and $a \in A$ define $\omega \cdot a \in A'$ by $\omega \cdot a(b) := \omega(ab)$ for $b \in A$. Now the space $\{\omega \cdot a : \omega \in A',\, a \in A\}$ is weak *-dense in A'. The cancellation property (2.1) of Δ_A shows that $\{(\omega \otimes \mathrm{id}_A)\Delta_A(A) :\ \omega \in A \text{ and } a \in A\}$ is norm dense in A; hence Δ_A is a faithful C*-action.

Theorem 4.4. *The C*-action γ of $\mathbb{G}$ on $\mathrm{C}_0(\widehat{G_1})$ is faithful if and only if β is non-trivial.*

Proof. If possible, assume that β is trivial. Then Proposition 3.2 gives

$$\gamma(\lambda(f)) = \Delta_{\mathrm{C}^*_{\mathrm{r}}(G_1)}(\lambda(f)) \otimes 1 \quad \text{for all } f \in \mathrm{C}_{\mathrm{c}}(G_1).$$

Here $\Delta_{\mathrm{C}^*_{\mathrm{r}}(G_1)}$ denotes the comultiplication on $\mathrm{C}^*_{\mathrm{r}}(G_1)$. By Example 4.3 we observe that the set $\{(\omega \otimes \mathrm{id}_A)\gamma(\lambda(f)) :\ \omega \in \mathrm{C}^*_{\mathrm{r}}(G_1)',\, f \in \mathrm{C}_{\mathrm{c}}(G_1)\}$ is strictly dense in $\mathcal{M}(\mathrm{C}^*_{\mathrm{r}}(G_1)$; hence γ is not faithful. Therefore, by contraposition we obtain that the faithfulness of γ implies the nontriviality of β.

Conversely, assume that β is non-trivial. By virtue of [1, Proposition 3.6] it is enough to show the set $\{(\omega \otimes \mathrm{id}_A)\gamma(\lambda(f)) :\ \omega \in \mathrm{C}^*_{\mathrm{r}}(G_1)'\}$ is norm dense in $\alpha(\mathrm{C}_0(G_2)) \cdot (\mathrm{C}^*_{\mathrm{r}}(G_1) \otimes 1)$.

Let K and K' be compact subsets of G_1 such that $\mu_1(K) \neq 0$ and $\mu_1(K') \neq 0$. Let χ and χ' be characteristic functions of K and K' respectively. For a given $f \in \mathrm{C}_{\mathrm{c}}(G_1)$ define

$$A_{\chi,\chi'} = (\omega_{\chi,\chi'} \otimes \mathrm{id}_A)\gamma(\lambda(f)).$$

Here $\omega_{\chi,\chi'}$ denotes the contraction with respect to χ_K and $\chi_{K'}$.

For all $\xi \in L^2(G_1 \times G_2)$ we get,

$$\begin{aligned} A_{\chi,\chi'}\xi(h,t) &= \int \overline{\chi(g)}\gamma(\lambda(f))(\chi'(g) \otimes \xi(h,t)\mathrm{d}\mu_1(g) \\ &= \iint \overline{\chi(g)}f(z)\chi'(g\beta_{\alpha_h(t)}(z))\xi(hz,t)\mathrm{d}\mu_1(g)\mathrm{d}\mu_1(z) \\ &= \int \mu_1(K \cap \beta_{\alpha_h(t)}(z)K')f(z)\xi(hz,t)\mathrm{d}\mu_1(z) \end{aligned}$$

Let $\{K_n\}_{n\in\mathbb{N}}$ be an increasing sequence of compact subsets of G_1 and growing up to the whole group G_1. Let χ_n denotes the characteristic function of K_n for all $n \in \mathbb{N}$. By dominated convergence theorem,

$$\lim_{n\to\infty} A_{\chi_n,\chi'}\xi(h,t) = \int \mu_1(\beta_{\alpha_h(t)}(z)K')f(z)\xi(hz,t)\mathrm{d}\mu_1(z).$$

Invariance of Haar measure μ_1 gives $\mu_1(\beta_{\alpha_h(t)}(z)K') = \mu_1(K')$ for all $h, z \in G_1$, $t \in G_2$. Therefore,

$$\lim_{n\to\infty} A_{\chi_n,\chi'}\xi(h,t) = \mu_1(K') \int f(z)\xi(hz,t)\mathrm{d}\mu_1(z) = \mu_1(K')(\lambda(f) \otimes 1)\xi(h,t).$$

Thus $A_{\chi_n,\chi}$ goes strictly to λ_{g_0} for some $g_0 \in G_1$. Therefore, for any $\eta, \eta' \in L^2(G_1)$, the operator $B_{\eta,\eta',g_0} := (\omega_{\eta,\eta'} \otimes \mathrm{id}_A)\gamma(\lambda(g_0))(\lambda_{g_0^{-1}} \otimes 1)$ is in the desired algebra.

Next we compute,

$$\begin{aligned} B_{\eta,\eta',g_0}\xi(h,t) &= \big((\omega_{\eta,\eta'}\otimes \mathrm{id}_A)\gamma(\lambda(g_0))\big)(\lambda_{g_0^{-1}}\otimes 1)\xi(h,t) \\ &= (\lambda_{g_0^{-1}}\otimes 1)\int \overline{\eta(g)}\eta'(g\beta_{\alpha_h(t)}(g_0))\xi(hg_0,t)\mathrm{d}\mu_1(g) \\ &= \int \overline{\eta(g)}\eta'(g\beta_{\alpha_h(t)}(g_0))\mathrm{d}\mu_1(g)\xi(h,t), \qquad \text{for } \xi\in L^2(G_1\times G_2). \end{aligned}$$

Since, β is non-trivial, varying η, η', g_0 and using Stone-Weierstrass theorem we get the set of functions $x\to\int\overline{\eta(g)}\eta'(g\beta_x(g_0))\mathrm{d}\mu_1(g)$ for $x\in G_2$ which is norm dense in $\mathrm{C}_0(G_2)$. Therefore, the set of operators B_{η,η',g_0} is norm dense in $\alpha(\mathrm{C}_0(G_2))$. □

4.2. **Isometry.** We recall the definition of isometric action from [5] for compact quantum groups. In analogy to this, it is natural to make the following definition of isometric action in the locally compact set-up:

Definition 4.5. Let γ be a C*-action of a locally compact quantum group $\mathbb{G}$ on $\mathrm{C}_0(X)$ where X is a smooth Riemannian (possibly non-compact) manifold with the Hodge-Laplacian $\mathcal{L}=-d^*d$, where d denotes the de-Rham differential operator. We say that γ is for every bounded linear functional ω on $M(Q)$, $(\mathrm{id}\otimes\omega)\circ\gamma$ maps $\mathrm{C}_{\mathrm{b}}^{\infty}(X)$ to itself and commutes with $\mathcal{L}$ on that subspace.

Just as in [6], it is easy to prove that for any isometric action γ and any smooth vector field χ on X, $f,\phi\in\mathrm{C}_{\mathrm{b}}^{\infty}(X)$, $(\chi\otimes\mathrm{id})(\gamma(f))$ and $\gamma(\phi)$ will commute.

Proposition 4.6. *Assume $\widehat{G_1}$ is a Lie group. Then the* C*-*action γ of $\mathbb{G}$ on $\mathrm{C}_0(\widehat{G_1})$ is isometric whenever either α or β is trivial.*

Proof. As noted before, the condition of isometry of γ implies the operators $(\delta_i\otimes\mathrm{id}_A)\gamma(\lambda_{g_1})$ and $\gamma(\lambda_{g_2})$ commute for all $g_1,g_2\in G_1$ and derivations δ_i on $\widehat{G_1}$.

Let $\xi\in L^2(G_1\times G_1\times G_2)$. Using Proposition 3.2, we compute

$$\begin{aligned} L &= (\delta_i\otimes\mathrm{id}_A)\gamma(\lambda_{g_1})\gamma(\lambda_{g_2})\xi(g,h,t) \\ &= \rho_i(\beta_{\alpha_h(t)}(g_1))\gamma(\lambda_{g_2})\xi\big(g\beta_{\alpha_h(t)}(g_1),hg_1,t\big) \\ &= \rho_i(\beta_{\alpha_h(t)}(g_1))\xi\big(g\beta_{\alpha_h(t)}(g_1)\beta_{\alpha_{hg_1}(t)}(g_2),hg_1g_2,t\big) \end{aligned}$$

Using (2.13) and commutativity of G_1 we get,

$$L=\rho_i(\beta_{\alpha_h(t)}(g_1))\xi\big(g\beta_{\alpha_h(t)}(g_1g_2),hg_1g_2,t\big).$$

A similar computation gives

$$R=\gamma(\lambda_{g_2})(\delta_i\otimes\mathrm{id}_A)\gamma(\lambda_{g_1})\xi(g,h,t)=\rho_i(\beta_{\alpha_{hg_2}(t)}(g_1))\xi\big(g\beta_{\alpha_h(t)}(g_1g_2),hg_1g_2,t\big).$$

Now $L=R$ for all $\xi\in L^2(G_1\times G_1\times G_2)$, $g_1,g_2\in G_1$, and $\rho_i\in\widehat{G_1}$, implies

$$\beta_{\alpha_h(t)}(g_1)=\beta_{\alpha_{hg_2}(t)}(g_1)\qquad\text{for all } g_1,g_2,h\in G_1,t\in G_2.$$

This is true if either of the actions α or β is trivial. □

We prove the main result of this section:

Theorem 4.7. *Assume $\widehat{G_1}$ is a Lie group and the* C*-*action γ of $\mathbb{G}$ on $\mathrm{C}_0(\widehat{G_1})$ is faithful and isometric. Then $\mathbb{G}$ is classical group.*

Proof. By Theorem 4.4, faithfulness of γ implies that β is non-trivial. On the other hand, γ is isometric; hence Proposition 4.6 forces α to be trivial. From (2.14) and using the fact that G_1 is abelian we get $M=L^{\infty}(\widehat{G_1}\times G_2)$. □

5. Examples

'$ax + b$' is the group of affine transformations of the real line $\mathbb{R}$. The natural action of $ax + b$ on $\mathbb{R}$ given by $x \mapsto ax + b$ for $a \in \mathbb{R} \setminus \{0\}$ and $b, x \in \mathbb{R}$ is faithful. We apply our results on two versions of quantum $ax + b$ group discussed in [15, Section 5]. Both of them are genuine non-compact, non-discrete, non-Kac quantum groups. We show that they act ergodically and faithfully on non-compact Riemannian manifolds. However none of these actions are not isometric.

5.1. **Baaj-Skandalis' ax+b group.** Assume $G_1 = G_2 = \mathbb{R} \setminus \{0\}$ and the group operation is the usual multiplication. Let $G = \{(a, b) \mid a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}\}$ with $(a, b)(c, d) = (ac, d + cb)$.

Define $i\colon G_1 \mapsto G$ and $j\colon G_2 \mapsto G$ by

$$i(g) := (g, g - 1), \qquad j(s) := (s, 0),$$

for all $g \in G_1$ and $s \in G_2$. This way (G_1, G_2) is a matched pair in the sense of Definition 2.11. Associated actions α and β are defined by

$$\alpha_g(s) = \frac{gs}{s(g-1)+1}, \qquad \beta_s(g) := s(g - 1) + 1.$$

for all $g, s \in \mathbb{R} \setminus \{0\}$ such that $(g - 1) \neq -s^{-1}$.

Associated bicrossed product $\mathbb{G}$ is the Baaj-Skandalis' quantum $ax + b$ group (see [15, Section 5.3]). By [15, Proposition 5.2 & 5.3], $\mathbb{G}$ is self dual, non-Kac, non-compact, non-discrete quantum group.

Proposition 5.1. *There is an ergodic, faithful and non-isometric* C*-*action of* $\mathbb{G}$ *on* $\mathrm{C}_0(\mathbb{R} \setminus \{0\})$.

Proof. Clearly, G_1 is abelian and $\widehat{G_1}$ is a Lie group with two connected components. Since β is non-trivial and $\mathbb{G}$ is a genuine quantum group, by Proposition 4.1, Theorem 4.4, and Theorem 4.7 the C*-action γ of $\mathbb{G}$ on $\mathrm{C}_0(\mathbb{R} \setminus \{0\})$ in Theorem 3.1 is ergodic, faithful and not isometric. □

5.2. **Split–Extension.** Assume $G_1 = \{(a, b) \mid a > 0, b \in \mathbb{R}\}$ with $(a, b)(c, d) := (ac, ad + \frac{b}{c})$ and $G_2 = (\mathbb{R}, +)$. Let K be the multiplicative group with two elements. Define $G = SL_2(\mathbb{R})/K$, and $i\colon G_1 \mapsto G$, $j\colon G_2 \mapsto G$ by

$$i(a, b) := \begin{pmatrix} a & b \\ 0 & \frac{1}{a} \end{pmatrix} \bmod K, \qquad j(x) := \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \bmod K.$$

This way (G_1, G_2) is a matched pair in the sense of Definition 2.11. Associated actions α and β are defined by

$$\alpha_{(a,b)}(x) := \frac{x}{a(a + bx)}, \qquad \beta_x(a, b) := \begin{cases} (a + bx, b) & \text{if } a + bx > 0, \\ (-a - bx, -b) & \text{if } a + bx < 0. \end{cases}$$

whenever $ax + b \neq 0$.

By [15, Proposition 5.5], associated bicrossed product $\mathbb{G}$ and its dual $\widehat{\mathbb{G}}$ are non-Kac, non-compact, and non-discrete quantum group. Also, $\widehat{\mathbb{G}}$ is not unimodular. Moreover, [15, Remark 5.6] shows that $\widehat{\mathbb{G}}$ is deformation of some generalised $ax + b$ group.

Proposition 5.2. *There is an ergodic, faithful and non-isometric* C*-*action of* $\widehat{\mathbb{G}}$ *on* $\mathrm{C}_0(\mathbb{R})$.

Proof. Clearly, G_2 is an abelian and $\widehat{G_2}$ is a connected Lie group. Recall that $\widehat{\mathbb{G}}$ is obtained by exchanging G_1 and α with G_2 and β. Since α is non-trivial, Proposition 4.1, Theorem 4.4, and Theorem 4.7 give the C*-action γ of $\widehat{\mathbb{G}}$ on $\mathrm{C}_0(\mathbb{R})$ in Theorem 3.1 is ergodic, faithful and not isometric. □

References

[1] Saad Baaj, Georges Skandalis, and Stefaan Vaes, *Non-semi-regular quantum groups coming from number theory*, Comm. Math. Phys. **235** (2003), no. 1, 139–167, doi: 10.1007/s00220-002-0780-6. MR 1969723

[2] Saad Baaj and Stefaan Vaes, *Double crossed products of locally compact quantum groups*, J. Inst. Math. Jussieu **4** (2005), no. 1, 135–173, doi: 10.1017/S1474748005000034. MR 2115071

[3] Vladimir Gershonovich Drinfel′d, *Quantum groups*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 1987, pp. 798–820. MR 934283

[4] Pavel Etingof and Chelsea Walton, *Semisimple Hopf actions on commutative domains*, Adv. Math. **251** (2014), 47–61, doi: 10.1016/j.aim.2013.10.008. MR 3130334

[5] Debashish Goswami, *Quantum group of isometries in classical and noncommutative geometry*, Comm. Math. Phys. **285** (2009), no. 1, 141–160, doi: 10.1007/s00220-008-0461-1. MR 2453592

[6] Debashish Goswami and Soumalya Joardar, *Rigidity of action of compact quantum groups on compact, connected manifolds*, eprint. arXiv:1309.1294.

[7] Huichi Huang, *Faithful compact quantum group actions on connected compact metrizable spaces*, J. Geom. Phys. **70** (2013), 232–236, doi: 10.1016/j.geomphys.2013.03.027. MR 3054297

[8] ———, *Invariant subsets under compact quantum group actions* (2013), eprint. arXiv:1210.5782v2.

[9] Michio Jimbo, *A q-difference analogue of $U(\mathfrak{g})$ and the Yang-Baxter equation*, Lett. Math. Phys. **10** (1985), no. 1, 63–69, doi: 10.1007/BF00704588. MR 797001

[10] Johan Kustermans and Stefaan Vaes, *Locally compact quantum groups*, Ann. Sci. École Norm. Sup. (4) **33** (2000), no. 6, 837–934, doi: 10.1016/S0012-9593(00)01055-7. MR 1832993

[11] ———, *Locally compact quantum groups in the von Neumann algebraic setting*, Math. Scand. **92** (2003), no. 1, 68–92, available at `http://www.mscand.dk/article.php?id=198`. MR 1951446

[12] Yuri Ivanovich Manin, *Quantum groups and noncommutative geometry*, Université de Montréal, Centre de Recherches Mathématiques, Montreal, QC, 1988. MR 1016381

[13] Tetsuya Masuda, Yoshiomi Nakagami, and Stanisław Lech Woronowicz, *A C^*-algebraic framework for quantum groups*, Internat. J. Math **14** (2003), no. 9, 903–1001, doi: 10.1142/S0129167X03002071. MR 2020804

[14] Ralf Meyer, Sutanu Roy, and Stanisław Lech Woronowicz, *Homomorphisms of quantum groups*, Münster J. Math. **5** (2012), 1–24, available at `http://nbn-resolving.de/urn:nbn:de:hbz:6-88399662599`. MR 3047623

[15] Stefaan Vaes and Leonid Vainerman, *Extensions of locally compact quantum groups and the bicrossed product construction*, Adv. Math. **175** (2003), no. 1, 1–101, doi: 10.1016/S0001-8708(02)00040-3. MR 1970242

[16] Stefaan Vaes and Alfons Van Daele, *Hopf C^*-algebras*, Proc. London Math. Soc. (3) **82** (2001), no. 2, 337–384, doi: 10.1112/S002461150101276X. MR 1806875

[17] Stanisław Lech Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), North-Holland, Amsterdam, 1998, pp. 845–884. MR 1616348

[18] ———, *From multiplicative unitaries to quantum groups*, Internat. J. Math. **7** (1996), no. 1, 127–149, doi: 10.1142/S0129167X96000086. MR 1369908

E-mail address: `goswamid@isical.ac.in`

Statistics and Mathematics Unit, Indian Statistical Institute, 203, B. T. Road, Kolkata 700108, India

E-mail address: `sutanu.roy@carleton.ca`

School of Mathematics and Statistics, Carleton University, 1125 Colonel By Drive, Ottawa, ON K1S 5B6, Canada